

Health Campaigns Have Reduced Armed Forces Veterans' Risk of Heart Attacks

Monday 12 May, 2014

Veterans who joined the Armed Forces in the 1960s and 1970s have suffered more heart attacks than their contemporaries who have never been in the military, according to a study by the University of Glasgow.

However, those who completed their service more recently are no longer at increased risk. The higher risk among older veterans was probably due to them being much more likely to smoke than their non-military contemporaries.

Recent health promotion campaigns adopted by the military have been successful in reducing smoking and the long term benefits are now being seen in younger veterans.

Researchers in the Institute of Health and Wellbeing looked at computerised and anonymised health records of 57,000 Scottish veterans who were born between 1945 and 1985 and served between 1960 and 2012. They compared their health records in later life with that of 173,000 non-service individuals to see if veterans were more likely to develop a range of health conditions including heart disease, cancer and mental health problems.

Their first report from the study, published in the American Journal of Epidemiology, shows that older veterans born in the 1940s and 1950s were more likely to suffer a heart attack than non-veterans of the same age. However, younger veterans have a similar risk to the wider population.

Dr Beverly Bergman, a PhD student and former British Army soldier and senior military doctor, conducted the research under the supervision of Professor Jill Pell. She said: "We wanted to do this research to improve understanding of veterans' health. We also want to provide important information to help local authorities, health boards, charities and the armed forces, to plan support services for veterans.

"There used to be a culture of heavy smoking and drinking in the army, but improved physical fitness and health promotion initiatives within the forces, which really came to the fore from the late 1970s onwards seem to have contributed to improved health, hence the difference we see between the two groups. It emphasises the importance of not smoking. Veterans who smoke can really make a difference by giving up.

"This is one of the few studies of its kind and is only really possible thanks to the excellent health records that Scotland has. It has shown us that we should not just be looking at veterans' mental health, as we now know they are at risk from heart attacks as well.

"We intend to look at other aspects of health to see whether veterans are more likely to suffer from certain issues than the general population."

Professor Jill Pell said: "Everyone is aware that the military are at greater risk of trauma than civilians. However, this study highlights that, in protecting their health, we mustn't overlook other lifestyle risk factors such as smoking that may continue to put their health at risk long after leaving the Service."

ENDS

Press Contact:

University of Glasgow press office 0044 141 330 3535 or media@glasgow.ac.uk

Related Sectors:

[Health](#) ::

Related Keywords:

[Armed Forces Veterans](#) ::

Scan Me:



Company Contact:

University of Glasgow

T. +44 (0)1413307126

E. nick.wade@glasgow.ac.uk ?

Additional Contact(s):

Nick Wade

[View Online](#)